




ELSEVIER

Contents lists available at ScienceDirect

Advances in Mathematics

www.elsevier.com/locate/aim



Weak differentiability for fractional maximal functions of general L^p functions on domains



João P.G. Ramos^a, Olli Saari^{a,*}, Julian Weigt^b

^a *Mathematical Institute, University of Bonn, Endenicher Allee 60, 53115, Bonn, Germany*

^b *Department of Mathematics and Systems Analysis, Aalto University, Finland*

ARTICLE INFO

Article history:

Received 24 November 2019

Received in revised form 26 March 2020

Accepted 1 April 2020

Available online xxxx

Communicated by Lars Olsen

MSC:

primary 42B25, 46E35

Keywords:

Maximal function

Sobolev space

Spherical means

Domains

ABSTRACT

Let $\Omega \subset \mathbb{R}^n$ be bounded domain. We prove under certain structural assumptions that the fractional maximal operator relative to Ω maps $L^p(\Omega) \rightarrow W^{1,p}(\Omega)$ for all $p > 1$, when the smoothness index $\alpha \geq 1$. In particular, the results are valid in the range $p \in (1, n/(n-1)]$ that was previously unknown. As an application, we prove an endpoint regularity result in the domain setting.

© 2020 Elsevier Inc. All rights reserved.

1. Introduction

Regularity of the Hardy–Littlewood maximal function of a Sobolev function was first studied in [13]. It was shown that the maximal operator preserves $W^{1,p}(\mathbb{R}^n)$ regularity for $p > 1$. This continues to hold true at the derivative level when $p = 1$ and $n = 1$

* Corresponding author.

E-mail addresses: jgramos@math.uni-bonn.de (J.P.G. Ramos), saari@math.uni-bonn.de (O. Saari), julian.weigt@aalto.fi (J. Weigt).

[21,16] and for radial functions [18]. Extending such a statement to more general Sobolev functions of several variables is a difficult open problem, which has inspired many results in related topics. For instance, slightly stronger bounds have been proved for maximal operators with more special convolution kernels (see [7], [3], [4] and [20]), the continuity of the mapping has been studied in [17] and [6], and a part of the techniques used for continuity, also relevant for the current paper, have been extended to $p = 1$ in [11].

Another aspect of the problem is the fractional endpoint question proposed by Carneiro and Madrid [5]. The fractional maximal function is given by

$$M_\alpha f(x) = \sup_{r>0} \frac{r^\alpha}{|B(x,r)|} \int_{B(x,r)} |f(y)| dy,$$

and it defines a bounded operator $L^p(\mathbb{R}^n) \rightarrow L^q(\mathbb{R}^n)$ when $q = np/(n-\alpha p)$, $0 < \alpha < n/p$ and $p > 1$. This boundedness fails at the endpoint $p = 1$, but the question about boundedness of ∇M_α from $W^{1,1}(\mathbb{R}^n)$ to $L^{n/(n-\alpha)}(\mathbb{R}^n)$ has not been answered so far for $\alpha < 1$ (see [19], [1] and [2] for related research and partial results). The case $\alpha \geq 1$ turned out to be very simple, and the reason can be traced back to the inequality

$$|\nabla M_\alpha f(x)| \leq c_{\alpha,n} M_{\alpha-1} f(x) \tag{1.1}$$

of Kinnunen and Saksman [15]. Carneiro and Madrid [5] noted that (1.1) together with the Gagliardo–Sobolev–Nirenberg inequality and the $L^p \rightarrow L^q$ bounds for the fractional maximal function imply the expected endpoint bound when $\alpha \geq 1$.

In the present paper, we study these problems in general open subsets of \mathbb{R}^n , which is a natural context for analysis from the point of view of potential theory and partial differential equations. Regularity of the local Hardy–Littlewood maximal function of a Sobolev function on an open $\Omega \subset \mathbb{R}^n$ was first studied by Kinnunen and Lindqvist [14], and a local variant of the inequality for the derivative of the fractional maximal function (1.1) was proved in [12]. This is our starting point, and for more thorough discussion of what was proved and what is unknown, we introduce some more notation.

If $\Omega \subset \mathbb{R}^n$ is an open set, the local fractional maximal function is defined as

$$M_\alpha^\Omega f(x) = \sup_{0 < r < \text{dist}(x, \Omega^c)} \frac{r^\alpha}{|B(x,r)|} \int_{B(x,r)} f(y) dy.$$

As the boundary of Ω restricts the choice of r in the definition, one cannot expect (1.1) to trivially carry over to the local setting. Indeed, such a pointwise inequality is false in general (Example 4.1 in [12]). On the other hand, if one adds a correction term involving the surface measure of the sphere to the right hand side of (1.1), one obtains

$$|\nabla M_\alpha^\Omega f(x)| \leq c_{\alpha,n} \left(M_{\alpha-1} f(x) + \sup_{r>0} |r^{\alpha-1} \sigma_r * f(x)| \right), \tag{1.2}$$

which is valid in all domains. This was used in [12] to prove that L^p functions with $p > n/(n - 1)$ large enough have $M_\alpha^\Omega f$ in a first order Sobolev class. The lower bound on p rules out functions too singular for an application of a spherical maximal function argument.

Our main theorem shows that under suitable assumptions on the domain Ω , the maximal function M_α^Ω maps $L^p(\Omega)$ into a first order Sobolev space for all $p > 1$.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^n$ be open, $n \geq 2$, $p > 1$ and $f \in L^p(\Omega)$. Then $M_\alpha^\Omega f$ is weakly differentiable and*

$$\|\nabla M_\alpha^\Omega f\|_{L^p(\Omega)} \leq C\|f\|_{L^p(\Omega)}$$

if any one of the following holds:

- (1) $\alpha > 1$ and Ω is bounded.
- (2) $\alpha = 1$ and Ω^c is convex.
- (3) $\alpha = 1$ and Ω is bounded and satisfies a uniform curvature bound in the sense of Section 2.2.
- (4) $\alpha = 1$ and $p > 1 + \frac{1}{n}$.

The constant C depends on the dimension, and in (1) and (3) it also depends on α and the domain.

Unlike [12], we are not able to prove an $L^p \rightarrow L^q$ smoothing effect on top of winning one derivative. However, our method does apply to singular functions in L^p spaces with $1 \leq p \leq n/(n - 1)$ where the argument in [12] fails to give any result. In particular, we have the following endpoint regularity result, which was previously out of reach.

Corollary 1.2. *Let $\Omega \subset \mathbb{R}^n$ be a Lipschitz domain. Then for all $f \in W^{1,1}(\Omega)$*

$$\|\nabla M_1^\Omega f\|_{L^{n/(n-1)}(\Omega)} \leq C\|f\|_{W^{1,1}(\Omega)}$$

where the constant C only depends on Ω and the dimension.

We briefly outline the proof of the main theorem. The maximal function on a domain behaves differently depending on whether the ball attaining the maximum touches the boundary or not. In case it does not, the local maximal function behaves like the global one, and the analysis is very similar. Otherwise it coincides with a linear averaging operator (2.3), which depends on the domain. These two parts are analyzed separately, and the main part of the proof is to establish L^p bounds for the derivative of (2.3). This leads to studying a domain dependent weighted spherical averaging operator (5.1).

Instead of resorting to maximal averages and the Bourgain–Stein theorem, an angular decomposition of the operator is carried out. The additional geometric information allows




Fig. 1. A set $P(y)$ and a tangent line.

instead to establish good L^1 bounds that can be interpolated with trivial L^∞ bounds in order to obtain a domination of (5.1) by a converging sum of L^p bounded operators. Improving the L^1 bound over what follows from the behavior of generic spherical means is crucial when aiming at L^p bounds for all $p > 1$. Such a conclusion cannot be drawn from mere polynomial decay of the Fourier transform of the weighted spherical measure in question, if no additional L^1 information is taken into account. Turning the focus from the Littlewood–Paley decomposition and L^2 methods to an angular decomposition and geometric estimates in L^1 is the leading insight of the proof.

The key idea in the L^1 estimates can be described as follows. Each domain Ω comes endowed with a family of sets (Fig. 1)

$$\{P(y) : y \in \Omega\}, \quad P(y) = \{x \in \Omega : y \in \partial B(x, \text{dist}(x, \Omega^c))\},$$

which can morally be used to dualize the spherical averaging operators (5.1) through Fubini’s theorem. The L^1 bounds for the constituents in the angular decomposition of the spherical averaging operator correspond to weighted integrals over the pieces of $P(y)$. If Ω is a ball, then the sets $P(y)$ are ellipsoids with foci at the center of the ball and at y . In the cases of the complement of a ball and a half-space, the $P(y)$ take the simple forms of hyperboloids and paraboloids. One cannot hope for as explicit descriptions as that in more general domains, but all $P(y)$ are boundaries of convex sets. This observation is used extensively in the proof.

The structure of the paper is as follows. In the first section, we introduce notation and some tools that will be helpful throughout the proof. The first sections are about differentiating the maximal function on so-called unconstrained points and proving the

weak differentiability of the maximal function conditionally to the L^p boundedness of the averaging operator (2.3). The rest of the paper is devoted to proving those L^p bounds by first computing a formula for the derivative and then carrying out the strategy sketched above. Finally, there is a concluding section with remarks on open problems and certain observations about the proof which might be of independent interest for future research.

2. Preliminaries

2.1. Notation

We let $n \geq 1$ denote the dimension. For a measurable set E , we let $|E|$ denote the n -dimensional Lebesgue measure. The k -dimensional Hausdorff measures are denoted by \mathcal{H}^k . An Euclidean ball with center $x \in \mathbb{R}^n$ and radius $r > 0$ is denoted by $B(x, r)$. A finite constant only depending on quantities that are not being kept track of is denoted by C . If $A \leq CB$ for such constant, we denote $A \lesssim B$ or write A is $\lesssim B$. We write $A \sim B$ if both $A \lesssim B$ and $B \lesssim A$ hold.

2.2. Domains

We always assume $\Omega \subset \mathbb{R}^n$ to be an open set, which we interchangeably call domain as the distinction obviously plays no role in this paper. We assume it to have non-empty complement. The distance function is denoted by $\delta(x) = \text{dist}(x, \Omega^c)$. As Ω^c is closed, there exists at least one $b_x \in \Omega^c$ so that $|x - b_x| = \delta(x)$. We reserve the notation b_x for such a point, which need not be unique unless Ω^c is convex. The distance function $\delta : \Omega \rightarrow [0, \infty)$ is always 1-Lipschitz. The gradient exists almost everywhere by Rademacher’s theorem, and it holds that

$$\nabla\delta(x) = \frac{x - b_x}{\delta(x)}. \tag{2.1}$$

This is because clearly the one sided directional derivative of $\delta(x)$ in the direction of $b_x - x$ always exists and is -1 . Where the gradient exists, we can use $|\nabla\delta(x)| \leq 1$ to conclude that the directional derivative in all directions orthogonal to $x - b_x$ must be zero.

A domain is said to satisfy a uniform curvature bound if there is an $R > 0$ so that for every point $x \in \Omega$ it holds

$$B\left(b_x + R\frac{x - b_x}{\delta(x)}, R\right) \subset \Omega.$$

All bounded C^2 domains satisfy this condition, but a domain satisfying a uniform curvature bound might be non-smooth and have inwards-pointing cusps. A domain with an interior ball condition need not satisfy the uniform curvature bound. An example of such a domain is $B(0, 1) \setminus (\{0\} \cup \{(0, \dots, 0, 2^k) : k \in \mathbb{Z}, k < 0\})$.

2.3. Function spaces on domains

Functions $f \in L^p(\Omega)$ are a priori only defined in the domain Ω , but we always extend them by zero to \mathbb{R}^n without additional comments. The Sobolev class $W^{1,p}(\Omega)$ consists of functions $f \in L^p(\Omega)$ such that $|\nabla f| \in L^p(\Omega)$. The weak derivatives are defined using test functions in $C_c^\infty(\Omega)$.

For the application of the main theorem to the endpoint regularity problem, we need a Sobolev embedding theorem for domains. One concrete case we can deal with is that of a Lipschitz domain.

Proposition 2.1 (Section 4.4 in [9]). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set so that $\partial\Omega$ is Lipschitz. Then for every $1 \leq p < \infty$ there exists a bounded extension operator*

$$E : W^{1,p}(\Omega) \rightarrow W^{1,p}(\mathbb{R}^n)$$

such that $\text{supp}(Ef) \subset B(x_0, 2 \text{diam}(\Omega))$ for some $x_0 \in \Omega$ and all $f \in W^{1,p}(\Omega)$.

By the boundary being Lipschitz, we mean that it can be covered by a finite number of open balls B_i so that for each i the domain $B_i \cap \Omega$ is the epigraph of a Lipschitz function.

The proposition together with the Gagliardo–Nirenberg–Sobolev inequality (see e.g. Section 4.5.1 in [9]) implies a rudimentary local Sobolev embedding

$$\|f\|_{L^{pn/(n-p)}(\Omega)} \leq C_{\Omega,p,n} \|f\|_{W^{1,p}(\Omega)} \tag{2.2}$$

valid for all $f \in W^{1,p}(\Omega)$ whenever Ω is a bounded open set with Lipschitz boundary. This is sufficient for our purposes.

2.4. Maximal function

For $\alpha \in [1, n)$, define the local fractional maximal function relative to Ω as

$$M_\alpha^\Omega f(x) = \sup_{0 < r < \delta(x)} r^\alpha \int_{B(x,r)} f(y) dy$$

whenever $f \in L^1_{\text{loc}}(\Omega)$. We omit the superscript when Ω is the whole \mathbb{R}^n . In addition, we define for $\alpha \in \mathbb{R}$ the auxiliary linear operator

$$A_\alpha f(x) = \delta(x)^\alpha \int_{B(z,\delta(x))} f(y) dy. \tag{2.3}$$

2.5. Constrained points

Let f be continuous. Fix $x \in \Omega$. Because the complement of Ω is non-empty, $\delta(x) < \infty$ and there exists a convergent sequence $r_j \in (0, \delta(x))$ with limit $r = \lim_{j \rightarrow \infty} r_j \in [0, \delta(x)]$ such that

$$M_\alpha^\Omega f(x) = \lim_{j \rightarrow \infty} r_j^\alpha \int_{B(x,r_j)} f(y) dy = r^\alpha \int_{B(x,r)} f(y) dy$$

if $r > 0$. If

$$M_\alpha^\Omega f(x) > \delta(x)^\alpha \int_{B(x,\delta(x))} f(y) dy,$$

the sequence r_j must be chosen so that $r < \delta(x)$, and the point x is said to be *unconstrained*. All other points are called *constrained*.

3. The unconstrained part

The local maximal function behaves similarly to the global one in the unconstrained set, and we reduce the differentiability question of the unconstrained part accordingly to that of the global maximal function. This is the content of the following proposition.

Proposition 3.1. *Let $p > 1$, $\alpha \geq 1$ and $f \in L^p_{loc}(\Omega)$ be continuous. The set U of the unconstrained points is open, the maximal function $M_\alpha^\Omega f$ is weakly differentiable in U , and the pointwise bound*

$$|\nabla M_\alpha^\Omega f(x)| \leq cM_{\alpha-1}f(x)$$

holds for a constant c only depending on the dimension and α whenever $x \in U$.

Proof. Consider the fractional average function

$$A(z, r) := r^\alpha \int_{B(z,r)} f(y) dy.$$

It is continuous in $(z, r) \in \Omega \times \mathbb{R}_+$. Fix now an unconstrained point x . By definition, there exists $\varepsilon > 0$ so that $M_\alpha^\Omega f(x) - A(x, \delta(x)) > \varepsilon$. Moreover, there exists $\gamma > 0$ so that $|(z, r) - (x, \delta(x))| < \gamma$, then $M_\alpha^\Omega f(x) - A(z, r) > \varepsilon/2$. Since $M_\alpha^\Omega f$ is lower semicontinuous, one can find for every z close enough to x a sequence $r_{z,j} \rightarrow r_z < \delta(x) - \gamma/2$ so that

$$M_\alpha^\Omega f(z) = \lim_{j \rightarrow \infty} r_{z,j}^\alpha \int_{B(z,r_{z,j})} f(y) dy.$$

In particular, there is an open neighborhood U_x of x so that for all $z \in U_x$

$$M_\alpha^\Omega f(z) = M_\alpha(1_{B(x,\delta(x))}f)(z).$$

By Theorem 3.1 in [15],

$$|\nabla M_\alpha^\Omega f(x)| \leq CM_{\alpha-1}f(x)$$

follows. \square

4. The full maximal function

Next we prove the differentiability of the local maximal function conditional to L^p bounds for the derivative of the averaging operator (2.3). This step morally follows from the lattice property of Sobolev functions, but as we only know the weak differentiability of $M_\alpha^\Omega f$ in the unconstrained set, some extra work is needed.

Lemma 4.1. *Let $p > 1$, $\alpha \geq 1$ and $\Omega \subset \mathbb{R}^n$ be such that ∇A_α and $M_{\alpha-1}$ are bounded $L^p(\Omega) \rightarrow L^p(\Omega)$. If $f \in L^p(\Omega)$, then the local fractional maximal function is weakly differentiable and*

$$\|\nabla M_\alpha^\Omega f\|_{L^p(\Omega)} \lesssim \|f\|_{L^p(\Omega)}.$$

Proof. Assume first that f is continuous and compactly supported. Following the arguments in [15], we infer that $M_\alpha^\Omega f$ can be seen as supremum over radii between a fixed upper and lower bound. The fractional averages are Lipschitz continuous with constants only depending on the radii, and hence their supremum is also Lipschitz. In particular, we know that $M_\alpha^\Omega f$ is continuous.

Denote by $g^+ = \max(g, 0)$ the positive part of a function g and write

$$M_\alpha^\Omega f = (M_\alpha^\Omega f - A_\alpha f)^+ + A_\alpha f.$$

By assumption, the second term admits the desired Sobolev bounds. To deal with the other term, let $\epsilon > 0$ and define

$$F_\epsilon(t) = \begin{cases} ((t - \epsilon)^2 + \epsilon^2)^{1/2} - \epsilon, & t > \epsilon \\ 0, & t \leq \epsilon. \end{cases}$$

These functions are of class $C^1(\mathbb{R})$ and converge pointwise to $t \mapsto (t)^+$ as $\epsilon \rightarrow 0$. Moreover, as $M_\alpha^\Omega f$ and $A_\alpha f$ are continuous, $E = \{x \in \Omega : F_\epsilon(M_\alpha^\Omega f(x) - A_\alpha f(x)) > 0\}$ has its closure contained in the open set of unconstrained points U . By Proposition 3.1, the assumption on A_α and the chain rule for Sobolev derivatives (4.2.2 in [9]), we obtain for all partial derivatives ∂_i

$$\partial_i F_\epsilon(M_\alpha^\Omega f - A_\alpha f) = (\partial_i M_\alpha^\Omega f - \partial_i A_\alpha f) F'_\epsilon(M_\alpha^\Omega f - A_\alpha f).$$

Taking a test function φ and computing

$$\int_\Omega F_\epsilon(M_\alpha^\Omega f - A_\alpha f) \partial_i \varphi \, dx = \int_\Omega (\partial_i M_\alpha^\Omega f - \partial_i A_\alpha f) F'_\epsilon(M_\alpha^\Omega f - A_\alpha f) \varphi \, dx,$$

we see that taking the limit $\epsilon \rightarrow 0$ proves the claim for continuous and compactly supported f .

To deal with the general case, let $f \in L^p(\Omega)$ and let f_j be continuous and compactly supported functions converging to f in L^p norm. By L^p continuity of the fractional maximal operator, $M_\alpha^\Omega f_j \rightarrow M_\alpha^\Omega f$ in L^p . As we have proved the following inequality

$$\|\nabla M_\alpha^\Omega f_j\|_{L^p(\Omega)} \lesssim \|f_j\|_{L^p(\Omega)},$$

for continuous functions f_j , the sequence $M_\alpha^\Omega f_j$ is bounded in $W^{1,p}(\Omega)$. We can extract a weakly convergent subsequence. By taking limits along this sequence and using the uniqueness of distributional limit, we conclude the proof for general $f \in L^p(\Omega)$. \square

As the main theorem is a direct consequence of the previous lemma, it remains to investigate the boundedness of the operator ∇A_α on $L^p(\Omega)$. The following sections are devoted to establishing the required L^p bounds when Ω is sufficiently well-behaved.

5. Constrained part

By a change of variables, we can write the averaging operator (2.3) as

$$A_\alpha f(x) = \delta(x)^\alpha \int_{B(0,1)} f(x + y\delta(x)) \, dy.$$

This operator is linear, and as we are aiming for L^p bounds, there is no loss of generality in restricting the attention to smooth functions. If x is a constrained point, then $M_\alpha^\Omega f(x) = A_\alpha f(x)$, which justifies our reference to A_α as the constrained part. Also, Lemma 4.1 showed that L^p bounds for the derivative of $A_\alpha f$ are enough to imply weak differentiability of the full maximal operator, so the maximal function does not play any role in what follows. A version of the following proposition was already proved in [12], but as we need a formula more precise than what they stated, we include the short proof for clarity.

Proposition 5.1. *Let $f \in C^\infty(\Omega)$. Then for almost every $x \in \Omega$*

$$|\nabla A_\alpha f(x)| \leq c_{n,\alpha} |A_{\alpha-1} f(x)| + c_n \delta(x)^{\alpha-1} \int_{\partial B(x,\delta(x))} \frac{|y - b_x|}{\delta(x)} f(y) d\mathcal{H}^{n-1}(y)$$

where $b_x \in \partial\Omega$ is a point such that $|b_x - x| = \delta(x)$.

Proof. Fix a point x . As $A_\alpha f(x) = \delta(x)^\alpha A_0 f(x)$, it holds that

$$\nabla A_\alpha f(x) = \alpha \delta(x)^{\alpha-1} A_0 f(x) \nabla \delta(x) + \delta(x)^\alpha (\nabla A_0 f)(x).$$

Since $|\nabla \delta(x)| \leq 1$ (cf. (2.1)), the first summand above is bounded by $A_{\alpha-1} f(x)$. Thus it suffices to analyze the gradient of $A_0 f$. Take the unit vector

$$e = \nabla A_0 f(x) / |\nabla A_0 f(x)|.$$

Then

$$\begin{aligned} |\nabla(A_0 f)(x)| &= (e \cdot \nabla) A_0 f(x) \\ &= \int_{B(0,1)} (e + y(e \cdot \nabla \delta(x))) \cdot \nabla f(x + \delta(x)y) dy \\ &= \frac{1}{\delta(x)} \int_{B(0,1)} \operatorname{div}_y ((e + y(e \cdot \nabla \delta(x))) f(x + \delta(x)y)) dy \\ &\quad - \frac{ne \cdot \nabla \delta(x)}{\delta(x)} \int_{B(0,1)} f(x + \delta(x)y) dy =: \text{I} + \text{II}. \end{aligned}$$

Since $|\nabla \delta(x)| \leq 1$, the contribution $\delta(x)^\alpha \cdot \text{II}$ is pointwise bounded by $nA_{\alpha-1} f$. To estimate the other term, we apply Gauss's theorem to obtain

$$\begin{aligned} \text{I} &= \frac{c_n}{\delta(x)} \int_{\partial B(0,1)} y \cdot (e + y(e \cdot \nabla \delta(x))) f(x + \delta(x)y) d\mathcal{H}^{n-1}(y) \\ &= \frac{c_n}{\delta(x)} \int_{\partial B(x,\delta(x))} \frac{(y - b_x) \cdot e}{\delta(x)} f(y) dy. \end{aligned}$$

So we reach the inequality

$$|\nabla A_\alpha f(x)| \leq \alpha n |A_{\alpha-1}(x)| + c_n \delta(x)^{\alpha-1} \int_{\partial B(x,\delta(x))} \frac{|y - b_x|}{\delta(x)} f(y) d\mathcal{H}^{n-1}(y),$$

which proves the claim. \square

Because $A_{\alpha-1}f(x) \leq M_{\alpha-1}^\Omega f(x)$, and $M_{\alpha-1}^\Omega$ satisfies the right $L^p \rightarrow L^q$ bounds, we have reduced the matter to understanding the weighted spherical average

$$B_\alpha f(x) := \delta(x)^{\alpha-1} \int_{B(x,\delta(x))} \frac{|y - b_x|}{\delta(x)} f(y) d\mathcal{H}^{n-1}(y) \tag{5.1}$$

on the right hand side of the conclusion of the previous proposition. The weight $|y - b_x|/\delta(x)$ measures the angle between $b_x - x$ and $y - x$ when $|y - b_x|/\delta(x)$ is small. We decompose the weighted spherical averaging operator according to the angle and location in the domain as follows. For $k \in \mathbb{Z}$, let

$$\Omega_k = \{x \in \Omega : 2^k \leq \delta(x) < 2^{k+1}\}$$

and for every point $x \in \Omega$ and integer $j \geq 0$

$$\omega_j(x) = \left\{ y \in \partial B(x, \delta(x)) : 2^{-j} < \frac{|y - b_x|}{\delta(x)} \leq 2^{-j+1} \right\}.$$

Define

$$S_j^k f(x) = 1_{\Omega_k}(x) \int_{\omega_j(x)} f(y) d\mathcal{H}^{n-1}(y).$$

Then

$$B_\alpha f(x) \lesssim \sum_{k \in \mathbb{Z}} \sum_{j=1}^\infty 2^{k(\alpha-n)-j} S_j^k f(x) \tag{5.2}$$

and it remains to prove bounds for S_j^k so that the right hand side sums up in L^p . This is done by interpolating bounds on L^∞ and L^1 .

Proposition 5.2. *Let Ω be any domain. It holds that $\|S_j^k\|_{L^\infty \rightarrow L^\infty} \lesssim 2^{(n-1)(k-j)}$, and consequently $\|\sum_k 2^{k(1-n)} S_j^k\|_{L^\infty \rightarrow L^\infty} \lesssim 2^{-(n-1)j}$.*

Proof. This follows from $\mathcal{H}^{n-1}(\omega_j(x)) \lesssim 2^{(n-1)(k-j)}$. \square

6. L^1 bounds

To prove L^1 bounds, we introduce some more notation. For each integer $j \geq 0$ and each point $y \in \Omega$, define

$$P_j(y) = \{x \in \Omega : y \in \omega_j(x)\}, \quad P(y) = \bigcup_{j=0}^\infty P_j(y). \tag{6.1}$$

In addition, let

$$A_j^k = \bigcup_{x \in \Omega_k} \omega_j(x). \tag{6.2}$$

Formally, certain weighted integrals over $P(y)$ give the adjoint operator of B_α . A naive change of order of integration is not justified in this case, but using the decomposition of B_α , we can make the idea precise. The following two propositions give effective description of $P(y)$ and provide a substitute for Fubini’s theorem.

Proposition 6.1. *Let Ω be an open set and let $y \in \Omega$. Then*

$$E(y) = \{x \in \Omega : |x - y| \leq \delta(x)\}$$

is closed and convex set such that

$$P(y) = \partial E(y).$$

For each $x \in P(y)$, the supporting hyperplane at x bisects the angle between $y - x$ and $b_x - x$ and is normal to $b_x - y$.

Proof. Recall that $P(y)$ consists of the points with $\{x \in \Omega : |y - x| = \delta(x)\}$. For $x \in P(y)$, it holds that

$$x + \epsilon \frac{b_x - x}{|b_x - x|} \in E(y)^c,$$

and it is easy to see $\partial E(y) = P(y)$. Consider the hyperplane

$$\{z \in \mathbb{R}^n : |z - b_x| = |z - y|\}.$$

It divides the space into two half spaces $H_1 = \{z : |z - b_x| < |z - y|\}$ and $H_2 = \{z : |z - b_x| \geq |z - y|\}$. If $x \in P(y)$, then $E(y) \subset H_2$ and $x \in H_2$. Thus ∂H_2 is a supporting hyperplane for $E(y)$ at x . As every boundary point of $E(y)$ has a supporting hyperplane, $E(y)$ is convex. The remaining assertions readily follow from the definition of ∂H_2 . \square

Proposition 6.2. *Let Ω be a domain, $j \geq 0$ and k integers and $f \geq 0$ a bounded continuous function on Ω . Then*

$$\int_{\Omega} S_j^k f(x) dx \lesssim 2^j \sum_{|j'-j| \leq 1, |k'-k| \leq 1} \int_{A_j^{k'}} f(y) \mathcal{H}^{n-1}(P_j^{k'}(y)) dy,$$

where we let $P_j^k(y) = P_j(y) \cap \Omega_k$.




Fig. 2. The construction to find x_1 .

Note that $y \in A_j^k$ if and only if $P_j^k(y) \neq \emptyset$.

Proof. The parameter k plays no role in the following computation, but is included in the statement for future reference. Let $\varphi \geq 0$ be a smooth function of one variable with compact support in $(0, 1)$ and $\|\varphi\|_{L^1(\mathbb{R})} = 1$. Denote the ϵ -dilation by $\varphi_\epsilon(t) = \epsilon^{-1}\varphi(t\epsilon^{-1})$. For any fixed x , we define the set of relevant directions

$$\omega_j^{\text{dir}}(x) = \delta(x)^{-1}(\omega_j(x) - x) \subset \partial B(0, 1).$$

As f is positive, the weak convergence

$$S_j^k f(x) = \int_{\omega_j(x)} f(y) d\mathcal{H}^{n-1}(y) = \lim_{\epsilon \rightarrow 0} \int_{x + \mathbb{R}\omega_j^{\text{dir}}(x)} f(y)\varphi_\epsilon(\delta(x) - |x - y|) dy$$

holds. Integrating over x , applying the dominated convergence theorem (this is justified, see the remark at the end of the argument), and using Fubini’s theorem, we obtain

$$\int_{\Omega_k} S_j f(x) dx \lesssim \int_{A_j^k} f(y) \left(\lim_{\epsilon \rightarrow 0} \frac{|\{x \in \Omega_k : y \in \omega_j^{\epsilon-}(x)\}|}{\epsilon} \right) dy \tag{6.3}$$

where the one-sided neighborhood is defined as (see Fig. 2)

$$\omega_j^{\epsilon-}(x) = x + \omega_j^{\text{dir}}(x)(\delta(x) - \epsilon, \delta(x)).$$

Next we estimate the limit expression in (6.3). As j and k are fixed, we can assume ϵ to be very small relative to them. Let $x \in \Omega_k$. Assume that $y \in \omega_j^{\epsilon-}(x)$. Then

$$-\epsilon < |y - x| - \delta(x) < 0 \tag{6.4}$$

and by definition x belongs to the interior of $E(y)$.

Set

$$e = \frac{b_x - x}{|b_x - x|}$$

and let $r \in (0, \delta(x))$ be such that $x + re \in P(y)$. Next we give an upper bound for r . Because $y \in \omega_j^{\epsilon^-}(x)$, it also holds that

$$\frac{y - x}{|y - x|} \in \omega_j^{\text{dir}}(x).$$

The mapping

$$g(\rho) := |y - (x + \rho e)| - \delta(x + \rho e) = |y - x - \rho e| - \delta(x) + \rho$$

is Lipschitz and hence absolutely continuous.

For all $\rho \geq 0$ we have the lower bound

$$\begin{aligned} g'(\rho) &= \partial_\rho[|y - (x + \rho e)| - \delta(x + \rho e)] = -e \cdot \frac{y - x - \rho e}{|y - x - \rho e|} + 1 \\ &= 1 - \cos \angle(b_x - x, y - x - \rho e) \\ &\geq 1 - \cos \angle(b_x - x, y - x) \\ &\gtrsim 2^{-2j} \end{aligned}$$

The last inequality is due to $y \in \omega_j^{\epsilon^-}(x)$. Recall that $g(0) \geq -\epsilon$ and $g(r) = 0$. Since g is absolutely continuous, we conclude

$$2^{-2j} r \lesssim \int_0^r g'(s) ds = g(r) - g(0) \leq \epsilon,$$

and

$$r \lesssim 2^{2j} \epsilon.$$

Denote $x_0 = x + re \in P(y)$. Consider the 2-plane containing x, y, b_x (and x_0). Its intersection with the convex body $E(y)$ provided by Proposition 6.1 is again a convex set E' in the plane. Let ℓ be its supporting line at x_0 . Then $x_0 \in P_j^{k'}(y)$ for some $j' \in \{j, j - 1\}, k' \in \{k, k - 1\}$ because

$$\begin{aligned} \angle(b_x - x_0, y - x_0) &\geq \angle(b_x - x, y - x) \geq 2^{-j} \\ \sin \angle(b_x - x_0, y - x_0) &\leq \frac{|b_x - y|}{\delta(x) - C2^{2j}\epsilon} = \frac{|b_x - y|\delta(x)^{-1}}{1 - C\delta(x)^{-1}2^{2j}\epsilon} \leq \sin 2^{-j+2} \end{aligned}$$

for ϵ small enough. By Proposition 6.1 this also means that $y - x_0$ makes an angle $\sim 2^{-j}$ with ℓ , and hence so does $x - x_0$. Let e' be the unit vector perpendicular to ℓ and $e' \cdot (y - x) < 0$. Then there is

$$s \lesssim |x - x_0| \sin 2^{-j} \lesssim 2^j \epsilon$$

so that $x + se' \in \ell$. Since $x \in E'(y)$ and ℓ intersects $E'(y)$ only in $\partial E'(y)$, there is $s' < s$ with $x_1 = x + s'e' \in \partial E'(y)$, which means

$$\text{dist}(x, P(y)) \lesssim 2^j \epsilon. \tag{6.5}$$

Recall that

$$\text{dist}(x, P_{j'}^{k'}(y)) \leq |x - x_0| \lesssim 2^{2j} \epsilon.$$

Let

$$N(\epsilon') = \bigcup_{j' \in \{j-1, j\}, k' \in \{k-1, k\}} \{x \in P(y) : \text{dist}(x, P_{j'}^{k'}(y)) \leq \epsilon'\}.$$

Then

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \frac{|\{x \in \Omega_k : y \in \omega_j^{\epsilon^-}(x) \leq \epsilon\}|}{\epsilon} &\leq \lim_{\epsilon' \rightarrow 0} \lim_{\epsilon \rightarrow 0} \frac{|\{x \in \Omega_k : \text{dist}(x, P(y)) \leq c_n 2^j \epsilon\} \cap N(\epsilon')|}{\epsilon} \\ &\lesssim \lim_{\epsilon' \rightarrow 0} 2^j \mathcal{H}^{n-1}(P(y) \cap N(\epsilon')) \\ &\leq \sum_{|j'-j| \leq 1, |k'-k| \leq 1} 2^j \mathcal{H}^{n-1}(P_{j'}^{k'}(y)), \end{aligned}$$

where the second inequality follows, for instance, by Theorem 3.2.39 in [10]. The integrable majorant of the sequence above that was needed for the application of the dominated convergence theorem before can be obtained by an application of the coarea formula. This completes the proof. \square

These two propositions are enough to conclude a general L^1 bound for the pieces S_j^k . This bound can be refined further, when additional regularity on the domain Ω is assumed.

Proposition 6.3. *Let Ω be an open set. Then $\|S_j^k\|_{L^1 \rightarrow L^1} \lesssim 2^{k(n-1)+j}$.*

Proof. If $x \in P(y) \cap \Omega_k$, then $|x - y| = \text{dist}(x, \Omega^c) \leq 2^{k+1}$. Hence $P(y) \cap \Omega_k \subset B(y, 2^{k+1})$. Recall that $P(y) = \partial E(y)$ and that $E(y)$ is convex. Thus $P(y) \cap \Omega_k \subset \partial(B(y, 2^{k+1}) \cap E(y))$ where $B(y, 2^{k+1}) \cap E(y)$ is convex. Since the perimeter of $B(y, 2^{k+1})$ dominates the perimeters of all convex sets with non-empty interior contained in it, we can conclude

$$\begin{aligned} \mathcal{H}^{n-1}(P_j^k(y)) &\leq \mathcal{H}^{n-1}(P(y) \cap B(y, 2^{k+1})) \leq \mathcal{H}^{n-1}(\partial(B(y, 2^{k+1}) \cap E(y))) \\ &\leq \mathcal{H}^{n-1}(\partial B(y, 2^{k+1})) \lesssim 2^{k(n-1)}. \end{aligned}$$

Now the claim follows from Proposition 6.2. \square

Remark 6.4. In case Ω is bounded and $\partial\Omega$ is C^2 smooth, the estimate for $\mathcal{H}^{n-1}(P_j^k(y))$ can be refined as follows. If $x \in P_j^k(y)$, then $|y - b_x| \leq \delta(x)2^{-j+1}$. This implies $\text{dist}(y, \partial\Omega) \leq \delta(x) \cdot 2^{-j+1}$ and further

$$|b_y - b_x| \leq |b_y - y| + |y - b_x| \leq 4\delta(x) \cdot 2^{-j}.$$

As the inward-pointing unit normal N_Ω at the boundary is well-defined and Lipschitz,

$$|N_\Omega(b_y) - N_\Omega(b_x)| \lesssim \text{diam}(\Omega)2^{-j}.$$

Because $N(b_z) = (z - b_z)/|z - b_z|$, this implies

$$\left| N_\Omega(b_y) - \frac{(x - y)}{\delta(x)} \right| \leq |N_\Omega(b_y) - N_\Omega(b_x)| + \frac{|y - b_x|}{\delta(x)} \lesssim \text{diam}(\Omega) \cdot 2^{-j}.$$

Therefore, all vectors $x - y$ with $y \in \omega_j(x)$ are within an angle $\sim \tilde{c}(\Omega) \cdot 2^{-j}$ of $N_\Omega(b_y)$. Hence the set $P_j^k(y)$ is contained in a cylinder of height $\sim 2^k$ and basis $\sim \tilde{c}(\Omega) \cdot 2^{k-j}$. By the inequality for perimeters of convex sets as in the proof of Proposition 6.3

$$\mathcal{H}^{n-1}(P_j^k(y)) \lesssim c(\Omega)2^k \cdot 2^{(k-j)(n-2)}.$$

This dependency on j is sharp even for very flat domains as can be seen by letting Ω be a smoothed out $B(0, 10) \cap \{x_1 \geq 0\}$ and $y = 2^{-j}e_1$ and $k \leq 0$.

However, as the estimate on $\mathcal{H}^{n-1}(P_j^k(y))$ is not the narrow gap of the proof of our main theorem, we do not pursue this aspect further.

The estimate $\mathcal{H}^{n-1}(P_j^k(y)) \lesssim 2^{k(n-1)}$ cannot be improved in general. If the boundary of the domain is a single point, the equality is achieved up to a constant. However, focusing on the whole $P_j(y)$ instead of single pieces $P_j^k(y)$, one can obtain a different estimate at cost of worsening the dependency on j . The following proposition is useful for small values of j , and it holds in very general domains.

Proposition 6.5. *Let Ω be an open set and $y \in \Omega$. Then*

$$\int_{P_j(y)} \frac{1}{\text{dist}(x, y)^{n-1}} d\mathcal{H}^{n-1}(x) \lesssim 2^j$$

with the constant independent of y . In particular,

$$\left\| \sum_k 2^{k(1-n)} S_j^k \right\|_{L^1 \rightarrow L^1} \lesssim 2^{2j}$$

Proof. We have

$$\int_{P_j(y)} \frac{1}{\text{dist}(x, y)^{n-1}} d\mathcal{H}^{n-1}(x) \lesssim \liminf_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \int_{\{x \in E(y)^c : \text{dist}(x, P_j(y)) \leq \epsilon\}} \frac{1}{\text{dist}(x, y)^{n-1}} dx.$$

Given any point $x \in P_j(y)$ and a line $l_x = \{y + t(x - y) : t \in \mathbb{R}\}$, we see that by Proposition 6.1 the line makes an angle $\sim 2^{-j}$ with $P_j(y)$, and hence

$$\mathcal{H}^1(l_x \cap \{z \in E(y)^c : \text{dist}(z, P_j(y)) \leq \epsilon\}) \lesssim 2^j \epsilon.$$

The first claimed bound for the integral follows immediately from passing to polar coordinates with origin at y .

To prove the second claim, note that by Proposition 6.2

$$\begin{aligned} \int_{\Omega} \sum_k 2^{k(1-n)} S_j^k f(x) dx &\lesssim 2^j \sum_{|j'-j| \leq 1} \int_{\Omega} f(y) \left(\sum_k 2^{k(1-n)} \mathcal{H}^{n-1}(P_{j'}^k(y)) \right) dy \\ &\lesssim 2^j \sum_{|j'-j| \leq 1} \int_{\Omega} f(y) \left(\int_{P_{j'}(y)} \frac{1}{\text{dist}(x, y)^{n-1}} d\mathcal{H}^{n-1}(x) \right) dy \\ &\lesssim 2^{2j} \|f\|_{L^1}, \end{aligned}$$

where the last step was an application of the first claim. \square

7. L^p bounds and geometry

To conclude bounds for the operator B_α , we have to sum up all the pieces in the decomposition. In order to make this work, one has to ensure that there is enough decay in j and k . Although the L^1 bounds do not sum up, interpolation with the better L^∞ bounds provides us with enough decay in the angle parameter j . If Ω is bounded, we can take advantage of the $L^p(\Omega)$ spaces being nested and use the decay in the scale parameter k near the boundary to complete the proof with no smoothness assumptions

on the boundary of the domain. This is possible only when we do not attempt to prove scalable estimates that would capture $L^p \rightarrow L^q$ smoothing beyond one derivative gain.

Theorem 7.1. *Let Ω be a bounded open set, $p, \alpha > 1$. Then*

$$\|B_\alpha\|_{L^p(\Omega) \rightarrow L^p(\Omega)} \lesssim \text{diam}(\Omega)^{\alpha-1}$$

where the implicit constant only depends on p, α and the dimension.

Proof. Let $S_j = \sum_k 2^{k(\alpha-n)} S_j^k$ so that $B_\alpha = \sum_j 2^{-j} S_j$. Then by Proposition 6.3

$$\begin{aligned} \|S_j\|_{L^1(\Omega) \rightarrow L^1(\Omega)} &\leq \sum_{k=-\infty}^{\log \text{diam}(\Omega)+1} 2^{k(\alpha-n)} \|S_j^k\|_{L^1(\Omega) \rightarrow L^1(\Omega)} \\ &\lesssim \sum_{k=-\infty}^{\log \text{diam}(\Omega)+1} 2^{k(\alpha-1)} 2^j \lesssim 2^j \text{diam}(\Omega)^{\alpha-1}. \end{aligned}$$

By Proposition 5.2

$$\|2^{-j} S_j\|_{L^\infty(\Omega) \rightarrow L^\infty(\Omega)} \lesssim 2^{-nj} \text{diam}(\Omega)^{\alpha-1}$$

and by interpolation we obtain

$$\|2^{-j} S_j\|_{L^p(\Omega) \rightarrow L^p(\Omega)} \lesssim 2^{-\frac{(p-1)n}{p} j} \text{diam}(\Omega)^{\alpha-1}.$$

As the exponent is negative, we can sum up in j to conclude the proof. \square

To deal with the critical case $\alpha = 1$ where our estimates have the correct scaling, we have to take into account finer properties of the boundary, as the estimation as rough as above leads to a logarithmic blow-up of the k -sum at the boundary.

Proposition 7.2. *Let Ω be an open set.*

- *If Ω satisfies the uniform curvature bound with R , then for all $y \in \Omega$ and $x \in P(y)$ with $\delta(x) \leq R$, it holds that*

$$\delta(x) \left(1 - \frac{\delta(x)}{R}\right) (1 - \cos \beta) \leq \text{dist}(y, \partial\Omega) \tag{7.1}$$

where $\beta = \angle(b_x - x, y - x)$.

- *If Ω^c is convex, then*

$$\delta(x) (1 - \cos \beta) \leq \text{dist}(y, \partial\Omega). \tag{7.2}$$




Fig. 3. The balls and points appearing in the proof of Proposition 7.2.

Proof. Take $x \in \Omega$ and $y \in \partial B(x, \delta(x))$ and let β be the angle between $b_x - x$ and $y - x$. Because Ω satisfies a uniform curvature bound, there is an $R > 0$ independent of x and y so that we can find a ball $B(z, R) \subset \Omega$ with $z = b_x + (x - b_x)R/\delta(x)$ so that $\overline{B(z, R)} \cap \partial\Omega = \{b_x\}$ (see Fig. 3). The Pythagorean identity reads

$$\begin{aligned} |z - y|^2 &= (\delta(x) \sin \beta)^2 + (R - \delta(x)(1 - \cos \beta))^2 \\ &= R^2 \left(1 - 2 \frac{\delta(x)}{R} \left(1 - \frac{\delta(x)}{R}\right) (1 - \cos \beta)\right) \\ &\leq R^2 \left(1 - \frac{\delta(x)}{R} \left(1 - \frac{\delta(x)}{R}\right) (1 - \cos \beta)\right)^2. \end{aligned}$$

Let w be the closest point to y in $\partial B(z, R)$. Since z, y and w are on the same line, we get

$$\begin{aligned} \text{dist}(y, \partial\Omega) &\geq |y - w| = |z - w| - |z - y| \\ &\geq R - R \left(1 - \frac{\delta(x)}{R} \left(1 - \frac{\delta(x)}{R}\right) (1 - \cos \beta)\right) \\ &= \delta(x) \left(1 - \frac{\delta(x)}{R}\right) (1 - \cos \beta) \end{aligned}$$

as claimed. If Ω^c is convex, then the uniform curvature bound is satisfied with $R = \infty$, whence the second claim follows. \square

Theorem 7.3. *Let Ω be an open set. Let $\alpha = 1$ and $p > 1$. Then*

- *If Ω is bounded and satisfies the uniform curvature bound, then*

$$\|B_1\|_{L^p(\Omega) \rightarrow L^p(\Omega)} \lesssim \log \left(\frac{\text{diam}(\Omega)}{R} + 1 \right)$$

where R is the radius from the uniform curvature bound.

- If Ω^c is convex, then

$$\|B_1\|_{L^p(\Omega) \rightarrow L^p(\Omega)} \lesssim 1$$

and the operator norm only depends on the dimension and p .

- If Ω is merely open, then

$$\|B_1\|_{L^p(\Omega) \rightarrow L^p(\Omega)} \lesssim 1$$

under the restriction $p > 1 + \frac{1}{n}$.

Proof. Proposition 6.3 implies

$$\int_{\Omega} 2^{k(1-n)-j} S_j^k f(x) dx \lesssim \int f(y) 1_{A_j^k}(y) dy. \tag{7.3}$$

Recall the definition (6.2). There are only $\sim \log(\text{diam}(\Omega)/R+1)$ values of k so that $R/8 \leq 2^k \leq 2 \text{diam}(\Omega)$. For k such that $2^{k+3} \leq R$, we can use the first item in Proposition 7.2 to see that for fixed y , the set $P_j^k(y)$ is non-empty only for k such that $2^{-2j+k} \lesssim \text{dist}(y, \partial\Omega)$. On the other hand, the upper bound

$$\text{dist}(y, \partial\Omega) \leq |y - b_x| \lesssim 2^{k-j}$$

is always valid, so $P_j^k(y)$ is non-empty only for $2^{-2j+k} \lesssim \text{dist}(y, \partial\Omega) \lesssim 2^{-j+k}$. Consequently,

$$A_j^k \subset \{y \in \Omega : 2^{-2j+k} \lesssim \text{dist}(y, \partial\Omega) \lesssim 2^{-j+k}\}.$$

For any y , there are only $\lesssim j$ values k such that the set above is non-empty, and hence by (7.3)

$$\left\| \sum_k 2^{k(1-n)-j} S_j^k \right\|_{L^1 \rightarrow L^1} \lesssim \log\left(\frac{\text{diam}(\Omega)}{R} + 1\right) + j.$$

Interpolation as in the proof of Theorem 7.1 implies the claim.

To prove the second item, just note that the convexity assumption on the complement means sending $R \rightarrow \infty$ so that $2^{k+3} \leq R$ always holds. To prove the third item, we study S_j as in the proof of Theorem 7.1 and replace the L^1 bound from Proposition 6.3 by that from Proposition 6.5. \square

Corollary 7.4. *Let Ω be a domain, $p > 1$ and $f \in L^p$. Then $A_\alpha f(x)$ from (2.3) is weakly differentiable and*

$$\|\nabla A_\alpha f\|_{L^p(\Omega)} \lesssim \|f\|_{L^p(\Omega)}$$

if any one of the following holds:

- $\alpha > 1$ and Ω is bounded.
- $\alpha = 1$ and Ω is bounded and satisfies a uniform curvature bound.
- $\alpha = 1$ and Ω^c is convex.
- $\alpha = 1$ and $p > 1 + \frac{1}{n}$

The constant depends on the domain, α and the dimension.

Proof. By linearity, it suffices to prove the norm inequality for smooth functions. By Proposition 5.1, it suffices to bound B_α from (5.1). This follows from Theorem 7.1 and Theorem 7.3 \square

Theorem 1.1 follows from Corollary 7.4 and Lemma 4.1.

8. Remarks

8.1. Role of the domain

It is not clear if the conditions on the domain in the hypothesis of Theorem 1.1 are necessary. One may ask if

$$\|\nabla M_1^\Omega\|_{L^p(\Omega) \rightarrow L^p(\Omega)} \lesssim 1$$

holds for all domains Ω and all $p > 1$. We are not aware of any counterexamples so far. Since M_0^Ω does satisfy an $L^p(\Omega)$ bound independent of the domain, the question is about the behavior of B_1 (see Theorem 7.3) in general domains. We point out that one avenue for improving the L^p bounds for B_1 could be to replace the strong L^1 bounds for S_j^k by weak type bounds in order to improve the operator norm bound with respect to j .

8.2. Endpoint regularity in domains

Corollary 1.2 follows from Theorem 1.1, since

$$\|\nabla M_1^\Omega f\|_{L^{n/(n-1)}(\Omega)} \lesssim \|f\|_{L^{n/(n-1)}(\Omega)} \lesssim \|f\|_{W^{1,1}(\Omega)}.$$

Here we used the main theorem and (2.2). The same observation was done by [5] to notice that the fractional endpoint regularity problem follows from inequality (1.1) as $\alpha \geq 1$ in the full space \mathbb{R}^n . The domain case was not known before as the inequality (1.1) should have been replaced by (1.2). This amounts to changing the Hardy–Littlewood maximal function to the spherical maximal function in the display above. That one is

not bounded in $L^{n/(n-1)}$, so the argument breaks down. However, using Theorem 1.1, we can complete the argument in certain domains Ω .

To the best of our knowledge, the fractional endpoint regularity problem has not been studied in domains before. It is hence natural to ask

Question 8.1. What must be assumed about an open set $\Omega \subset \mathbb{R}^n$ so that

$$\|\nabla M_\alpha^\Omega f\|_{L^{\alpha/(n-\alpha)}(\Omega)} \lesssim \|f\|_{W^{1,1}(\Omega)}$$

for $\alpha \in (0, 1]$?

Our main theorem gives some information on the case $\alpha = 1$, but the remaining values of α remain open. The values $\alpha > 1$ can be dealt with using a spherical maximal function argument with no additional assumptions. The remaining values of α are probably way harder to handle as the endpoint regularity question is completely open even in the full space.

Finally, we remark that the techniques used to get results for smooth kernels as in [2] are insensitive to the ambient domain, because one does not use precise information about the maximizing radius. The arguments there only rely on sublinearity of maximal functions. Hence a $W^{1,1}$ variant of Theorem 1.1 in [2] easily extends to the domain setting. Indeed, fixing $\alpha \in (0, 1)$, letting Ω be any Sobolev extension domain, $\Omega_\epsilon = \{x \in \Omega : \text{dist}(x, \Omega^c) \leq \epsilon\}$ and m a local maximal function with kernel compactly supported and smooth enough as in [2], one can invoke Theorem 3 in Section 5.8.2 in [8] to reduce the problem to proving

$$\lim_{\epsilon \rightarrow 0} \sup_{h \in B(0, \epsilon/2)} \int_{\Omega_\epsilon} \left| \frac{mf(x+h) - mf(x)}{|h|} \right|^{\frac{n}{n-\alpha}} dx \lesssim \|f\|_{W^{1,1}(\Omega)}^{\frac{n}{n-\alpha}}.$$

As $f \in W^{1,1}(\Omega)$ coincides with its extension $Ef \in W^{1,1}(\mathbb{R}^n)$ for all $x \in \Omega$, the integral on the left hand side can be controlled by a maximal multiplier as in [2] acting on $Ef(\cdot + h) - Ef(\cdot)$. Then the claim follows from Theorem 3.1 in [2] and the assumed boundedness of $E : W^{1,1}(\Omega) \rightarrow W^{1,1}(\mathbb{R}^n)$.

8.3. Smoothing for cube maximal functions

An equally interesting variant of the local fractional maximal function is the one defined by taking averages over cubes instead of balls

$$M_\alpha^{\Omega, \text{cube}} f(x) = \sup_{r>0, Q(x,r) \subset \Omega} r^\alpha \int_{Q(x,r)} f(y) dy.$$

As the faces of the cubes are completely flat, there are no L^p bounds for the maximal function

$$\sup_{r>0} \int_{\partial Q(x,r)} f(y) d\mathcal{H}^{n-1}(y), \tag{8.1}$$

and this was singled out as the principal reason why the methods in [12] do not extend to the case of cubical fractional maximal function.

Although we avoid the use quantities of the type (8.1), our proof is also inapplicable to the cubical case. There are two obvious obstructions:

- Let Ω be the upper half-plane. Take $\delta > 0$ and define f as the characteristic function of $[-\delta, \delta] \times [0, \delta^s]$ for some $s \geq 1$. Varying s and sending $\delta \rightarrow 0$, we see that

$$\|B_1 f\|_{L^p} \lesssim \|f\|_{L^p}$$

cannot hold for any $p < \infty$.

- As a detail in the proof, one can note that the analogues of the sets $P(y)$ from (6.1) defined relative to cubes might have full measure. The role of curvature, or lack of it, manifests in the 2^j factor in the statement of Proposition (6.2).

On the other hand, it seems that the problems with the cubical maximal function are not only a matter of lack of curvature. As the remarks above show, there are domains where averages over flat surfaces cause problems. However, if the geometry of the domain is very special, this kind of phenomena can be ruled out. The following observation gives an example.

Proposition 8.2. *Let $\Omega = \{(x, y) \in \mathbb{R}^2 : x < y\}$. Then*

$$\|\nabla M_\alpha^{\Omega, cube} f\|_{L^p} \lesssim \|f\|_{L^p}$$

for all $f \in L^p$.

Sketch of proof. The reduction to the cubical analogue of (5.2) follows by the lines of the spherical proof. Then it suffices to note that the decomposition in j and k is unnecessary, and an L^p bound for $p > 1$ follows by Minkowski’s inequality and a change of variables. \square

The exact behavior of the cubical local fractional maximal function in more general domains remains an interesting open problem.

8.4. Scalable estimates

The method of the proof of Theorem 1.1 forced us to prove $L^p \rightarrow W^{1,p}$ estimates for the derivative of the fractional maximal function. Such estimates can only hold true in bounded domains or for $\alpha = 1$, and in bounded domains they are weaker than the

expected $L^p \rightarrow \dot{W}^{1, \frac{np}{p-(\alpha-1)}}$ estimates, only known for $p > n/(n-1)$ by [12]. We do not pursue this possible improvement direction here, although we believe it to be an interesting open problem.

Acknowledgments

We would like to thank Juha Kinnunen for turning our attention to the error term in [12] and Martí Prats for comments and corrections to the previous version of the paper. Part of the research was done when the second author was visiting the Nonlinear PDE group at Aalto University, which he wishes to thank for its hospitality. JPGR was supported by the Deutscher Akademischer Austauschdienst (DAAD). OS was supported by Hausdorff Center for Mathematics DFG EXC 2047 and the project DFG SFB 1060. JW was supported by the Väisälä Foundation and the Academy of Finland.

References

- [1] D. Beltran, J. Madrid, Endpoint Sobolev continuity of the fractional maximal function in higher dimensions, *Int. Math. Res. Not. IMRN* (2019), <https://doi.org/10.1093/imrn/rnz281>, arXiv:1906.00496, 2017.
- [2] D. Beltran, J.P. Ramos, O. Saari, Regularity of fractional maximal functions through Fourier multipliers, *J. Funct. Anal.* 276 (6) (2019) 1875–1892.
- [3] E. Carneiro, R. Finder, M. Sousa, On the variation of maximal operators of convolution type II, *Rev. Mat. Iberoam.* 34 (2) (2018) 739–766.
- [4] E. Carneiro, C. González-Riquelme, Gradient bounds for radial maximal functions, arXiv:1906.01487, 2019.
- [5] E. Carneiro, J. Madrid, Derivative bounds for fractional maximal functions, *Trans. Am. Math. Soc.* 369 (6) (2017) 4063–4092.
- [6] E. Carneiro, J. Madrid, L.B. Pierce, Endpoint Sobolev and BV continuity for maximal operators, *J. Funct. Anal.* 273 (10) (2017) 3262–3294.
- [7] E. Carneiro, B.F. Svaiter, On the variation of maximal operators of convolution type, *J. Funct. Anal.* 265 (5) (2013) 837–865.
- [8] L.C. Evans, *Partial Differential Equations*, second edition, Graduate Studies in Mathematics, vol. 19, American Mathematical Society, Providence, RI, 2010.
- [9] L.C. Evans, R.F. Gariepy, *Measure Theory and Fine Properties of Functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1992.
- [10] H. Federer, *Geometric Measure Theory*, Die Grundlehren der mathematischen Wissenschaften, vol. 153, Springer-Verlag, New York Inc., New York, 1969.
- [11] P. Hajlasz, J. Malý, On approximate differentiability of the maximal function, *Proc. Am. Math. Soc.* 138 (1) (2010) 165–174.
- [12] T. Heikkinen, J. Kinnunen, J. Korvenpää, H. Tuominen, Regularity of the local fractional maximal function, *Ark. Mat.* 53 (1) (2015) 127–154.
- [13] J. Kinnunen, The Hardy-Littlewood maximal function of a Sobolev function, *Isr. J. Math.* 100 (1997) 117–124.
- [14] J. Kinnunen, P. Lindqvist, The derivative of the maximal function, *J. Reine Angew. Math.* 503 (1998) 161–167.
- [15] J. Kinnunen, E. Saksman, Regularity of the fractional maximal function, *Bull. Lond. Math. Soc.* 35 (4) (2003) 529–535.
- [16] O. Kurka, On the variation of the Hardy-Littlewood maximal function, *Ann. Acad. Sci. Fenn., Math.* 40 (1) (2015) 109–133.
- [17] H. Luiro, Continuity of the maximal operator in Sobolev spaces, *Proc. Am. Math. Soc.* 135 (1) (2007) 243–251.
- [18] H. Luiro, The variation of the maximal function of a radial function, *Ark. Mat.* 56 (1) (2018) 147–161.

- [19] H. Luiro, J. Madrid, The variation of the fractional maximal function of a radial function, *Int. Math. Res. Not. IMRN* (17) (2019) 5284–5298.
- [20] C. Pérez, T. Picon, O. Saari, M. Sousa, Regularity of maximal functions on Hardy-Sobolev spaces, *Bull. Lond. Math. Soc.* 50 (6) (2018) 1007–1015.
- [21] H. Tanaka, A remark on the derivative of the one-dimensional Hardy-Littlewood maximal function, *Bull. Aust. Math. Soc.* 65 (2) (2002) 253–258.