

Higher Dimensional Techniques for the Regularity of Maximal Functions

Julian Weigt

University of Warwick

18.05.2023

Introduction: Background

For $f : \mathbb{R}^n \rightarrow \mathbb{R}$ the centered Hardy-Littlewood maximal function is defined by

$$M^c f(x) = \sup_{r>0} f_{B(x,r)} \quad \text{with} \quad f_{B(x,r)} = \frac{1}{\mathcal{L}(B(x,r))} \int_{B(x,r)} |f|.$$

Theorem (Hardy-Littlewood maximal function theorem)

$$\|M^c f\|_{L^p(\mathbb{R}^n)} \leq C_{n,p} \|f\|_{L^p(\mathbb{R}^n)}$$

if and only if $p > 1$.

Introduction: Background

Theorem (Juha Kinnunen (1997))

For $p > 1$ we have

$$\|\nabla M^c f\|_{L^p(\mathbb{R}^n)} \leq C_{n,p} \|\nabla f\|_{L^p(\mathbb{R}^n)}$$

Proof: For $e \in \mathbb{R}^n$ by the sublinearity of M^c

$$\begin{aligned} \partial_e M^c f(x) &\sim \frac{M^c f(x + he) - M^c f(x)}{h} \\ &\leq \frac{M^c(f(\cdot + he) - f)(x)}{h} \\ &= M^c\left(\frac{f(\cdot + he) - f}{h}\right)(x) \sim M^c(\partial_e f)(x) \end{aligned}$$

By the Hardy-Littlewood maximal function theorem for $p > 1$

$$\|\nabla M^c f\|_{L^p(\mathbb{R}^n)} \lesssim \|M^c(|\nabla f|)\|_{L^p(\mathbb{R}^n)} \lesssim \|\nabla f\|_{L^p(\mathbb{R}^n)}$$

Question (Hajlasz and Onninen 2004)

Is it true that

$$\|\nabla M^c f\|_{L^1(\mathbb{R}^n)} \leq C_n \|\nabla f\|_{L^1(\mathbb{R}^n)}?$$

Uncentered Hardy-Littlewood maximal function

$$\tilde{M}f(x) = \sup_{B \ni x} f_B.$$

Endpoint question by Hajlasz and Onninen is interesting for \tilde{M} and other maximal operators.

Introduction: In one dimension

Theorem (Tanaka 2002, Aldaz and Pérez Lázaro 2007)


For $f : \mathbb{R} \rightarrow \mathbb{R}$ we have

$$\|\nabla \tilde{M}f\|_1 \leq \|\nabla f\|_1$$

In one dimension:

$$\|\nabla f\|_1 = \sup_{x_1 < x_2 < \dots} \sum_i |f(x_{i+1}) - f(x_i)| = \text{var } f$$

and $\tilde{M}f$ is convex on connected components of $\{x \in \mathbb{R} : \tilde{M}f(x) > f(x)\}$.



$$\text{var}_{[x_0, x_1]} \tilde{M}f \leq \text{var}_{[x_0, x_1]} f$$

Introduction: The fractional maximal function

For $0 < \alpha < n$ the centered fractional Hardy-Littlewood maximal function is

$$M_{\alpha}^c f(x) = \sup_{r>0} r^{\alpha} f_{B(x,r)}.$$

Corresponding Hardy-Littlewood theorem

$$\|M_{\alpha} f\|_{L^{p_{\alpha}}(\mathbb{R}^n)} \leq C_{n,\alpha,p} \|f\|_{L^p(\mathbb{R}^n)}$$

with $p_{\alpha} = \frac{pn}{n-\alpha p} > p$ if and only if $p > 1$. Corresponding regularity bound

$$\|\nabla M_{\alpha} f\|_{L^{p_{\alpha}}(\mathbb{R}^n)} \leq C_{n,\alpha,p} \|\nabla f\|_{L^p(\mathbb{R}^n)},$$

proven for $p > 1$.

Introduction: The fractional maximal function

Theorem (Kinnunen and Saksman 2003)

For $\alpha \geq 1$

$$|\nabla M_\alpha^c f(x)| \lesssim |M_{\alpha-1}^c f(x)|.$$

For $\alpha \geq 1$ we have $1_\alpha = \left(\frac{n}{n-1}\right)_{\alpha-1}$ and therefore

$$\begin{aligned} \|\nabla M_\alpha^c f\|_{L^{1_\alpha}(\mathbb{R}^n)} &\lesssim \|M_{\alpha-1}^c f\|_{L^{1_\alpha}(\mathbb{R}^n)} \lesssim \|f\|_{L^{\frac{n}{n-1}}(\mathbb{R}^n)} \\ &\lesssim \|\nabla f\|_{L^1(\mathbb{R}^n)}. \end{aligned}$$

- What about $0 < \alpha < 1$?
- Same result for \tilde{M}_α .

Introduction: Past progress

$n = 1$

[Tanaka 2002, Aldaz
+Pérez Lázaro 2007]

block decreasing f

[Aldaz+Pérez Lázaro 2009]

centered M , $n = 1$

[Kurka 2015]

radial f

[Luiro 2018]

fractional:

$n = 1$

[Beltran + Madrid 2016]

$1 \leq \alpha$

[Kinnunen + Saksman 2003
Carneiro + Madrid 2016]

radial f

[Luiro + Madrid 2017]

lacunary

[Beltran + Ramos + Saari 2018]

$n = 1$, radial for centered f

[Beltran + Madrid 2019]

more related bounds, bounds on other maximal operators, such as local, . . . , for example: Continuity of $f \mapsto \nabla Mf$ on $W^{1,1}(\mathbb{R}^n) \rightarrow L^1(\mathbb{R}^n)$, stronger than boundedness.

Introduction: New results

We prove the endpoint regularity bound for the maximal function for

- characteristic f
- dyadic maximal operator
- fractional maximal operator
- cube maximal operator

Introduction: Proof ingredients

Coarea formula

$$\|\nabla f\|_{L^1(\mathbb{R}^n)} = \int_{\mathbb{R}} \mathcal{H}^{n-1}(\partial\{x \in \mathbb{R}^n : f(x) > \lambda\}) d\lambda$$

Superlevel sets

$$\{Mf > \lambda\} = \{x \in \mathbb{R}^n : Mf(x) > \lambda\} = \bigcup \{B : f_B > \lambda\}$$

for *uncentered* maximal operators.

Decomposition of the boundary

Denote

$$\mathcal{B}_\lambda^< = \{B : f_B > \lambda, \mathcal{L}(B \cap \{f > \lambda\}) < 2^{-n-1}\mathcal{L}(B)\}$$

and $\mathcal{B}_\lambda^>$ accordingly.

Introduction: Proof ingredients

1 relative isoperimetric inequality:

$$\min\{\mathcal{L}(Q \cap E), \mathcal{L}(Q \setminus E)\}^{n-1} \lesssim \mathcal{H}^{n-1}(Q \cap \partial E)^n.$$

2 **Vitaly covering** and similar: general balls \rightarrow separated balls

3 **Besicovitch covering for boundary**

4 **superlevelset estimate:** $f < 0$ on most of $B \Rightarrow$ most mass of f lies far above f_B

	isoperimetric, Vitali	boundary Besicovitch	superlevel
dyadic char. f.	x		
char. f.	x	x	
dyadic	x		x
fractional	x		x
cube	x	x	x

Proof: Reformulation and decomposition

We have

$$\{Mf > \lambda\} = \bigcup \mathcal{B}_\lambda^< \cup \bigcup \mathcal{B}_\lambda^{\geq}.$$

Since $\{f > \lambda\} \subset \{Mf > \lambda\}$ we have

$$\partial\{Mf > \lambda\} \subset (\partial\{Mf > \lambda\} \setminus \overline{\{f > \lambda\}}) \cup \partial\{f > \lambda\}.$$

We conclude

Decomposition


$$\begin{aligned} \int_{\mathbb{R}^d} |\nabla Mf| &\leq \int_0^\infty \mathcal{H}^{n-1}(\partial \bigcup \mathcal{B}_\lambda^{\geq} \setminus \overline{\{f > \lambda\}}) \, d\lambda \\ &\quad + \int_0^\infty \mathcal{H}^{n-1}(\partial \bigcup \mathcal{B}_\lambda^<) \, d\lambda \end{aligned}$$

Proof: High density case $\mathcal{B}_\lambda^{\geq}$

Proposition

For Q, E with $\mathcal{L}(Q \cap E) \geq 2^{-n-1} \mathcal{L}(Q)$ we have

$$\mathcal{H}^{n-1}(\partial Q \setminus \bar{E}) \lesssim \mathcal{H}^{n-1}(Q \cap \partial E)$$



dyadic maximal operator


$$M^d f(x) = \sup_{\text{dyadic } Q, Q \ni x} f_Q.$$

$$\begin{aligned} \mathcal{H}^{n-1}(\partial \bigcup Q_\lambda^\geq \setminus \overline{\{f > \lambda\}}) &\leq \sum_{Q \in Q_\lambda^\geq} \mathcal{H}^{n-1}(\partial Q \setminus \overline{\{f > \lambda\}}) \\ &\lesssim \sum_{Q \in Q_\lambda^\geq} \mathcal{H}^{n-1}(Q \cap \partial\{f > \lambda\}) \\ &\leq \mathcal{H}^{n-1}(\partial\{f > \lambda\}) \end{aligned}$$

Proposition

For a set B of balls B with $\mathcal{L}(B \cap E) \geq 2^{-n-1} \mathcal{L}(B)$ we have

$$\mathcal{H}^{n-1}(\partial \bigcup B \setminus \overline{E}) \lesssim \mathcal{H}^{n-1}(\bigcup B \cap \partial E).$$



Proof: Low density case $\mathcal{B}_\lambda^<$, dyadic

$$\int_{\mathbb{R}} \mathcal{H}^{n-1}(\partial \cup \mathcal{Q}_\lambda^<) d\lambda \leq \sum_{Q \text{ dyadic}} (f_Q - \lambda_Q) \mathcal{H}^{n-1}(\partial Q)$$

with

$$\mathcal{L}(Q \cap \{f > \lambda_Q\}) = 2^{-n-1} \mathcal{L}(Q)$$

Proposition

$$(f_Q - \lambda_Q) \mathcal{L}(Q) \lesssim \int_{f_Q}^{\infty} \mathcal{L}(Q \cap \{f > \lambda\}) d\lambda$$

Proof: Low density case $\mathcal{B}_\lambda^<$, dyadic

$$\int_{\mathbb{R}} \mathcal{H}^{n-1}(\partial \cup \mathcal{Q}_\lambda^<) d\lambda \leq \sum_{Q \text{ dyadic}} (f_Q - \lambda_Q) \mathcal{H}^{n-1}(\partial Q)$$

with


$$\mathcal{L}(Q \cap \{f > \lambda_Q\}) = 2^{-n-1} \mathcal{L}(Q)$$

Proposition

$$(f_Q - \lambda_Q) \mathcal{L}(Q) \lesssim \int_{f_Q}^{\infty} \sum_{P \subsetneq Q: \bar{\lambda}_P < \lambda < f_P} \mathcal{L}(P \cap \{f > \lambda\}) d\lambda$$

where P is maximal above $\bar{\lambda}_P$ and

$$\mathcal{L}(P \cap \{f > \bar{\lambda}_P\}) = 2^{-1} \mathcal{L}(P)$$



Proof: Low density case $\mathcal{B}_\lambda^<$, dyadic

Combining, we obtain

$$\begin{aligned} \int_{\mathbb{R}} \mathcal{H}^{n-1}(\partial \bigcup Q_\lambda^<) d\lambda \\ \lesssim \int_{\mathbb{R}} \sum_Q \sum_{\text{dyadic } P \subseteq Q: \bar{\lambda}_P < \lambda < f_P} \frac{\mathcal{L}(P \cap \{f > \lambda\})}{l(Q)} d\lambda \end{aligned}$$

- 1 change the order of summation
- 2 convergence of the geometric sum
- 3 apply the relative isoperimetric inequality to P .
- 4 coarea formula to recover $\|\nabla f\|_1$

Proof: Low density case $\mathcal{B}_\lambda^<$, fractional

$1 \leq \alpha$ [Kinnunen + Saksman, Carneiro + Madrid]

$$\|\nabla M_\alpha f\|_{\frac{n}{n-\alpha}} \lesssim \|M_{\alpha-1} f\|_{\frac{n}{n-\alpha}} \lesssim \|f\|_{\frac{n}{n-1}} \lesssim \|\nabla f\|_1.$$

$0 < \alpha$

$$\|\nabla M_\alpha f\|_{\frac{n}{n-\alpha}} \lesssim \|M_{\alpha,-1} f\|_{\frac{n}{n-\alpha}} \lesssim \|\nabla f\|_1,$$

$M_{\alpha,-1}$ replacement for $M_{\alpha-1}$ if $0 < \alpha < 1$.

Can bound $M_{\alpha,-1} f$ both centered and uncentered

- using low density arguments from the dyadic proof
- extra flexibility coming from $\alpha > 0$, allowing for rough Vitali covering arguments

Proof: Low density case $\mathcal{B}_\lambda^<$, general cubes

cube maximal function

$$Mf(x) = \sup_{\text{cube } Q, Q \ni x} f_Q.$$

We reduce to almost dyadic cubes, using

Proposition (Vitali/Besicovitch for perimeter)

For any (finite) set of cubes \mathcal{Q} there is a subset $\mathcal{S} \subset \mathcal{Q}$ of disjoint cubes such that

$$\mathcal{H}^{n-1}(\partial \cup \mathcal{Q}) \lesssim \sum_{S \in \mathcal{S}} \mathcal{H}^{n-1}(\partial S).$$

Uncentered HL $\tilde{M}f$ (balls)?

- All arguments work
- except low density bound $(f_B - \lambda_B)\mathcal{L}(B) \lesssim?$

Thank you