

1. Find a bounded Lebesgue integrable function on $[0, 1]$ which is not Riemann integrable and prove these properties. /3

Hint: A function of the form 1_E will do.

2. Let $E \subset \mathbb{R}^d$ be $\mathcal{L}$ -measurable. Show, that there exist compact sets $K_1, K_2, \dots \subset E$ with $\mathcal{L}(K_n) \rightarrow \mathcal{L}(E)$. /1

3. Let $f : \Omega \rightarrow [0, \infty]$ be μ -measurable.

- (a) Show, that the area below the graph, $\{(x, t) \in \Omega \times \mathbb{R} : 0 \leq t < f(x)\}$, is $\mu \times \mathcal{L}$ -measurable. *Hint: It can be written as a countable union of "rectangles" $E \times [0, t)$ for certain measurable $E \subset \Omega$ and $t \in \mathbb{R}$.* /3

- (b) Show the **layer cake formula** / **Cavalieri's principle**, /2

$$\int_{\Omega} f \, d\mu = \int_0^{\infty} \mu(\{f > \lambda\}) \, d\lambda.$$

Hint: Fubini

- (c) For $1 \leq p < \infty$ prove the L^p -variant /3

$$\int_{\Omega} f^p \, d\mu = p \int_0^{\infty} \lambda^{p-1} \mu(\{f > \lambda\}) \, d\lambda$$

and conclude **Chebyshev's inequality**:

$$\mu(\{f > \lambda\}) \leq \frac{1}{\lambda^p} \int_{\Omega} f^p \, d\mu.$$

- (d) The space of functions f for which $\sup_{\lambda > 0} \lambda^p \mu(\{|f| > \lambda\}) < \infty$ is called $L^{p,\infty}(\Omega, \mu)$, or **weak- $L^p(\Omega, \mu)$** . Part (c) implies $L^{p,\infty}(\Omega, \mu) \subset L^p(\Omega, \mu)$. /2

Show that the inclusion is strict for $\Omega = \mathbb{R}$ and $\mu = \mathcal{L}$, i.e. that there exists a measurable $f : \mathbb{R} \rightarrow [0, \infty]$ with $\sup_{\lambda > 0} \lambda^p \mathcal{L}(\{f > \lambda\}) < \infty$ but $\|f\|_{L^p(\mathbb{R}, \mathcal{L})} = \infty$.

4. For $\lambda > 0$ and $E \subset \mathbb{R}^d$ denote $\lambda E = \{\lambda x : x \in E\}$.

- (a) Show, that /2

$$\mathcal{L}_*(\lambda E) = \lambda^d \mathcal{L}_*(E).$$

- (b) Show, that E is Lebesgue measurable if and only if λE is. /2